

Cuadro IV.3.1 PROMEDIO DE CONCENTRACIONES DEL CONTAMINANTE
PM-10 EN LA CIUDAD DE PANAMÁ, POR ESTACIÓN, SEGÚN MES:
AÑOS 2000-02

| Mes | PM-10 (ug/m ³) | | | | | | | | | | | |
|-----------------|----------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | 2000 | | | | 2001 | | | | 2002 | | | |
| | Ba | UP | HP | SM | Ba | UP | HP | SM | Ba | UP | HP | SM |
| Promedio..... | 33.4 | 62.0 | 43.8 | 87.2 | 38.6 | 66.2 | 47.9 | 92.4 | 25.8 | 53.2 | 36.9 | 72.2 |
| Enero..... | 31.0 | 60.8 | 46.4 | 71.6 | 49.7 | 76.4 | 49.7 | 90.2 | 35.7 | 57.3 | 28.4 | 42.3 |
| Febrero..... | 40.8 | 33.5 | 34.9 | 48.6 | 51.5 | 73.6 | 52.4 | 70.4 | 21.8 | 49.4 | 41.3 | 47.3 |
| Marzo..... | 28.5 | 78.8 | 40.0 | 55.2 | 35.6 | 75.3 | 42.1 | 82.1 | 28.4 | 56.4 | 43.4 | 55.3 |
| Abril..... | ... | 69.1 | 89.7 | 63.1 | 37.2 | 64.9 | 45.3 | 72.7 | 24.0 | 57.5 | ... | ... |
| Mayo..... | 30.2 | 58.9 | 59.2 | 122.4 | 30.2 | 46.3 | 47.8 | 98.5 | 25.3 | 45.3 | ... | ... |
| Junio..... | 41.6 | 67.0 | 40.7 | 103.7 | 27.2 | 75.6 | 54.7 | 120.9 | 22.4 | 60.0 | 28.0 | 86.2 |
| Julio..... | 26.0 | 73.4 | 22.6 | 98.6 | ... | 51.0 | 43.5 | 111.8 | 20.6 | 44.3 | 32.5 | 91.3 |
| Agosto..... | 30.1 | 43.1 | 33.4 | 89.9 | ... | ... | ... | ... | 28.1 | 55.3 | 47.8 | 110.8 |
| Septiembre..... | 33.7 | 50.3 | 35.1 | 96.5 | ... | ... | ... | ... | 25.7 | 53.0 | 37.0 | 72.0 |
| Octubre..... | 31.8 | 73.7 | 39.5 | 97.3 | ... | ... | ... | ... | ... | ... | ... | ... |
| Noviembre..... | 33.5 | 62.0 | 40.1 | 99.2 | ... | ... | ... | ... | ... | ... | ... | ... |
| Diciembre..... | 40.5 | 73.4 | 44.3 | 100.1 | ... | ... | ... | ... | ... | ... | ... | ... |

Estaciones: Balboa(BA), Universidad de Panamá (UP); Hipódromo Presidente Remón (HP);
San Miguelito (SM).

Fuente: Universidad de Panamá, Instituto Especializado de Análisis (IEA).

